



Finitness of the basic intersection cohomology of a Killing foliation

Martintxo Saralegi-Aranguren, Robert Wolak

► To cite this version:

Martintxo Saralegi-Aranguren, Robert Wolak. Finitness of the basic intersection cohomology of a Killing foliation. *Mathematische Zeitschrift*, 2012, 272, pp.443-457. 10.1007/s00209-011-0942-3 . hal-00443657

HAL Id: hal-00443657

<https://univ-artois.hal.science/hal-00443657>

Submitted on 1 Jan 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Finiteness of the basic intersection cohomology of a Killing foliation

Martintxo Saralegi-Aranguren*

Université d'Artois

Robert Wolak†

Uniwersytet Jagiellonski

January 3, 2010

Abstract

We prove that the basic intersection cohomology $H_{\overline{p}}^*(M/\mathcal{F})$, where \mathcal{F} is the singular foliation determined by an isometric action of a Lie group G on the compact manifold M , is finite dimensional.

This paper deals with an action $\Phi: G \times M \rightarrow M$ of a Lie group on a compact manifold preserving a riemannian metric on it. The orbits of this action define a singular foliation \mathcal{F} on M . Putting together the orbits of the same dimension we get a stratification of M . This structure is still very regular. The foliation \mathcal{F} is in fact a conical foliation and we can define the basic intersection cohomology $H_{\overline{p}}^*(M/\mathcal{F})$ (cf. [10]). This invariant becomes the basic cohomology $H^*(M/\mathcal{F})$ when the action Φ is almost free, and the intersection cohomology $H_{\overline{p}}^*(M/G)$ when the Lie group G is compact.

The aim of this work is to prove that this cohomology $H_{\overline{p}}^*(M/\mathcal{F})$ is finite dimensional. This result generalizes [3] (almost free case), [11] (abelian case) and [10] (compact case).

The paper is organized as follows. In Section 1 we present the foliation \mathcal{F} . The basic intersection cohomology $H_{\overline{p}}^*(M/\mathcal{F})$ associated to this foliation is studied in Section two. Twisted products are studied in Section 3. The finiteness of $H_{\overline{p}}^*(M/\mathcal{F})$ is proved in Section 4.

In the sequel M is a connected, second countable, Hausdorff, without boundary and smooth (of class C^∞) manifold of dimension m . All the maps are considered smooth unless something else is indicated.

1 Killing foliations determined by isometric actions.

We study in this work the foliations induced by isometric actions: the *Killing foliations*. These foliations are examples of the conical foliations for which the basic intersection cohomology has been defined (see [10, 11]). We present this geometrical framework in this section.

*Univ Lille Nord de France F-59 000 LILLE, FRANCE. UArtois, Laboratoire de Mathématiques de Lens EA 2462. Fédération CNRS Nord-Pas-de-Calais FR 2956. Faculté des Sciences Jean Perrin. Rue Jean Souvraz, S.P. 18. F-62 300 LENS, FRANCE. saralegi@euler.univ-artois.fr.

†Instytut Matematyki. Uniwersytet Jagiellonski. Stanisława Łojasiewicza 6, 30 348 KRAKOW, POLAND. robert.wolak@im.uj.edu.pl. Partially supported by the KBN grant 2 PO3A 021 25.

1.1 Killing foliations. A smooth action $\Phi: G \times M \rightarrow M$ of a Lie group G on a manifold M is a *isometric action* when there exists a riemannian metric μ on M preserved by G .

The connected components of the orbits of the action Φ determine a partition \mathcal{F} on M . In fact, this partition is a singular riemannian foliation that we shall call *Killing foliation* (cf. [7]). Notice that \mathcal{F} is also a conical foliation in the sense of [10, 11]. Classifying the points of M following the dimension of the leaves of \mathcal{F} one gets the *stratification* $\mathbf{S}_{\mathcal{F}}$ of \mathcal{F} . It is determined by the equivalence relation $x \sim y \Leftrightarrow \dim G_x = \dim G_y$. The elements of $\mathbf{S}_{\mathcal{F}}$ are called *strata*.

In the particular case where the closure of G in the isometry group of (M, μ) is a compact Lie group¹ we shall say that the action Φ is a *tame action*. In fact, a smooth action $\Phi: G \times M \rightarrow M$ is tame if and only if it extends to a smooth action $\Phi: K \times M \rightarrow M$ where K is a compact Lie group containing G (cf. [6]). The group K is not unique, but we always can choose K in such a way that G is dense in K . We shall say that K is a *tamer group*. Here the strata of $\mathbf{S}_{\mathcal{F}}$ are K -invariant closed submanifolds of M .

Since the aim of this work is the study of \mathcal{F} and not the action Φ itself, we can consider that the Lie group G is connected. Let us see that.

Proposition 1.1.1 *Let $\Phi: G \times M \rightarrow M$ is a tame action. Let G_0 be the connected component of G containing the unity element. The Killing foliation defined by the restriction $\Phi: G_0 \times M \rightarrow M$ is also \mathcal{F} .*

Proof. The partition \mathcal{F} is defined by this equivalence relation:

$$x \sim y \iff \exists \text{ continuous path } \alpha: [0, 1] \rightarrow G(x) \text{ such that } \alpha(0) = x \text{ and } \alpha(1) = y.$$

Since the map $\Delta: G \rightarrow G(x)$, defined by $\Delta(g) = \Phi(g, x) = g \cdot x$, is a submersion (see for example [2]) then

$$x \sim y \iff \exists \text{ continuous path } \beta: [0, 1] \rightarrow G \text{ such that } \beta(0) = e \text{ and } \beta(1) \cdot x = y,$$

and by definition of G_0

$$x \sim y \iff \exists \text{ continuous path } \beta: [0, 1] \rightarrow G_0 \text{ such that } \beta(0) = e \text{ and } \beta(1) \cdot x = y.$$

This gives the result. ♣

When G is connected, the tamer group K has richer properties.

Proposition 1.1.2 *Let G be a connected Lie subgroup of a compact Lie group K . If G is dense in K then $G \triangleleft K$ and the quotient group K/G is commutative.*

Proof. The Lie algebra \mathfrak{g} is Ad_G -invariant and hence, by density, Ad_K -invariant. Then \mathfrak{g} is an ideal of \mathfrak{k} . The connectedness of G gives that G is a normal subgroup of K . Since Ad_G acts trivially on $\mathfrak{k}/\mathfrak{g}$, Ad_K acts trivially, too. Therefore, $\mathfrak{k}/\mathfrak{g}$ is abelian (see for example [8, pag. 628]). ♣

1.2 Particular tame actions. A *trio* is a triple (K, G, H) , with K is a compact Lie group, G a normal subgroup of K and H a closed subgroup of K . We present now some tame actions associated to a trio (K, G, H) . They are going to be intensively used in this work. First of all we need some definitions.

- The action $\Phi_l: K \times K \rightarrow K$ is defined by $\Phi_l(g, k) = g \cdot k$. For each element u of the Lie algebra \mathfrak{k} of K , we shall write X^u the associated (right invariant) vector field. It is defined by $X^u(k) = T_e R_k(u)$ where $R_k: K \rightarrow K$ is given by $R_k(\ell) = \ell \cdot k$.

¹This is always the case when the manifold M is a compact.

- The action $\Phi_r: K \times K \rightarrow K$ is defined by $\Phi_r(g, k) = k \cdot g^{-1}$. For each element $u \in \mathfrak{k}$ of K , we shall write X_u the associated (left invariant) vector field. It is defined by $X_u(k) = -T_e L_k(u)$ where $L_k: K \rightarrow K$ is given by $L_k(\ell) = k \cdot \ell$.
- The action $\Psi: K \times K/H \rightarrow K/H$ is defined by $\Psi(g, kH) = (g \cdot k)H$. For each element $u \in \mathfrak{k}$, we shall write Y_u the associated vector field. Since the canonical projection $\pi: K \rightarrow K/H$ is a K -equivariant map, then we have $\pi_* X^u = Y_u$ for each $u \in \mathfrak{k}$.
- The action $\Gamma: H \times H \rightarrow H$ is defined by $\Gamma(g, h) = g \cdot h$. For each element u of the Lie algebra \mathfrak{h} of H we write Z^u the associated (right invariant) vector field.

The associated actions we are going to use are the following.

(a) *The restriction $\Phi_l: G \times K \rightarrow K$, which induces the regular Killing foliation \mathcal{K} .*

(b) *The restriction $\Phi_r: G \times K \rightarrow K$, which induces the regular Killing foliation \mathcal{K} .*

Since $G \triangleleft K$, the foliation \mathcal{K} is determined by the family of vector fields $\{X^u / u \in \mathfrak{g}\}$, where \mathfrak{g} is the Lie algebra of G , and also by the family $\{X_u / u \in \mathfrak{g}\}$. The orbits $G(k) = Gk = kG$ have the same dimension $\dim G$.

(c) *The restriction $\Psi: G \times K/H \rightarrow K/H$, which induces the regular Killing foliation \mathcal{D} .*

The foliation \mathcal{D} is determined by the family of vector fields $\{Y_u / u \in \mathfrak{g}\}$. The orbits $G(kH)$ have the same dimension $\dim G - \dim(G \cap H)$.

(d) *The restriction $\Gamma: (G \cap H) \times H \rightarrow H$, which induces the regular Killing foliation \mathcal{C} .*

The foliation \mathcal{C} is determined by the family of vector fields $\{Z^u / u \in \mathfrak{g} \cap \mathfrak{h}\}$. The orbits $(G \cap H)(k)$ have the same dimension $\dim(G \cap H)$.

(e) *The restriction $\Phi_r: GH \times K \rightarrow K$, which induces the regular Killing foliation \mathcal{E} .*

Notice that GH is a Lie group since G is normal in K . The foliation \mathcal{E} is, in fact, determined by the vector fields $\{X_u / u \in \mathfrak{g} + \mathfrak{h}\}$. The orbits $(GH)(k)$ have the same dimension $\dim G + \dim H - \dim(G \cap H)$.

1.3 Twisted product. In order to prove the finiteness of the basic intersection cohomology we decompose the manifold in a finite number of simpler pieces. These are the twisted products we introduce now.

We fix a trio (K, G, H) and a smooth action $\Theta: H \times N \rightarrow N$ of H on the manifold N . The *twisted product* is the quotient $K \times_H N$ of $K \times N$ by the equivalence relation $(k, z) \sim (k \cdot h^{-1}, \Theta(h, z) = h \cdot z)$. The element of $K \times_H N$ corresponding to $(k, z) \in K \times N$ is denoted by $\langle k, z \rangle$. This manifold is endowed with the tame action

$$\Phi: G \times (K \times_H N) \longrightarrow (K \times_H N),$$

defined by $\Phi(g, \langle k, z \rangle) = \langle g \cdot k, z \rangle$. We denote by \mathcal{W} the induced Killing foliation.

We also use the following tame action, namely, the restriction

$$\Theta: (G \cap H) \times N \rightarrow N$$

whose induced Killing foliation is denoted by \mathcal{N} .

The canonical projection $\Pi: K \times N \rightarrow K \times_H N$ relates the involved foliations as follows:

(a) $\Pi_*(\mathcal{K} \times \mathcal{I}) = \mathcal{W}$, where \mathcal{I} is the pointwise foliation (since the map Π is G -equivariant).

(b) $\mathcal{S}_{\mathcal{W}} = \{\Pi(K \times S) / S \in \mathcal{S}_{\mathcal{N}}\} = \Pi(\{K\} \times \mathcal{S}_{\mathcal{N}})$ (since $G_{\langle k, z \rangle} = k(G \cap H)_z k^{-1}$).

2 Basic Intersection cohomology

In this section we recall the definition of the basic intersection² cohomology and we present the main properties we are going to use in this work. For the rest of this section, we fix a conical foliation \mathcal{F} defined on a manifold M . The associated stratification is $\mathbf{S}_{\mathcal{F}}$. The regular stratum of is denoted by $R_{\mathcal{F}}$. We shall write $m = \dim M$, $r = \dim \mathcal{F}$ and $s = m - r = \text{codim}_M \mathcal{F}$.

We are going to deal with differential forms on a product $(\text{manifold}) \times [0, 1]^p$, they are restrictions of differential forms defined on $(\text{manifold}) \times]-1, 1[^p$.

2.1 Perverse forms. Recall that a *conical chart* is a foliated diffeomorphism $\varphi: (\mathbb{R}^{m-n-1} \times c\mathbb{S}^n, \mathcal{H} \times c\mathcal{G}) \rightarrow (U, \mathcal{F}_U)$ where $(\mathbb{R}^{m-n-1}, \mathcal{H})$ is a simple foliation and $(\mathbb{S}^n, \mathcal{G})$ is a conical foliation without 0-dimensional leaves. We also shall denote this chart by (U, φ, S) where S is the stratum of $\mathbf{S}_{\mathcal{F}}$ verifying $\varphi(\mathbb{R}^{m-n-1} \times \{\vartheta\}) = U \cap S$.

The differential complex $\Pi_{\mathcal{F}}^*(M \times [0, 1]^p)$ of *perverse forms* of $M \times [0, 1]^p$ is introduced by induction on depth $\mathbf{S}_{\mathcal{F}}$. When this depth is 0 then

$$\Pi_{\mathcal{F}}^*(M \times [0, 1]^p) = \Omega^*(M \times [0, 1]^p).$$

Consider now the generic case. A perverse form of $M \times [0, 1]^p$ is first of all a differential form $\omega \in \Omega^*(R_{\mathcal{F}} \times [0, 1]^p)$ such that,

$$\begin{cases} \text{the pull-back} & (\varphi \times \mathbb{I}_{[0, 1]^p})^* \omega \in \Omega^*(\mathbb{R}^{m-n-1} \times R_{\mathcal{G}} \times]0, 1[\times [0, 1]^p) \\ \text{extends to} & \omega_{\varphi} \in \Pi_{\mathcal{H} \times c\mathcal{G}}^*(\mathbb{R}^{m-n-1} \times \mathbb{S}^n \times [0, 1]^{p+1}) \end{cases}$$

for any conical chart (U, φ) , where \mathbb{I}_{\bullet} stands for the identity map. Notice that $\Omega^*(M)$ is included on $\Pi_{\mathcal{F}}^*(M)$ ³.

2.2 Perverse degree. The amount of transversality of a perverse form $\omega \in \Pi_{\mathcal{F}}^*(M)$ with respect to a singular stratum $S \in \mathbf{S}_{\mathcal{F}}$ is measured by the perverse degree $\|\omega\|_S$. We recall here the definition of local perverse degree $\|\omega\|_U \in \{-\infty\} \cup \mathbb{N}$ of ω relatively to a conical chart (U, φ, S) :

1. $\|\omega\|_U = -\infty$ when $\omega_{\varphi} \equiv 0$ on $\mathbb{R}^{m-n-1} \times R_{\mathcal{G}} \times \{0\}$,
2. $\|\omega\|_U \leq p$, with $p \in \mathbb{N}$, when $\omega_{\varphi}(v_0, \dots, v_p, -) \equiv 0$ where the vectors $\{v_0, \dots, v_p\}$ are tangent to the fibers of $P_{\varphi}: \mathbb{R}^{m-n-1} \times R_{\mathcal{G}} \times \{0\} \rightarrow U \cap S$ ⁴.

This number does not depend on the choice of the conical chart (cf. [11, Proposition 1.3.1]). Finally, we define the *perverse degree* $\|\omega\|_S$ by

$$\|\omega\|_S = \sup \left\{ \|\omega\|_U / (U, \varphi, S) \text{ conical chart} \right\}.$$

The perverse degree of $\omega \in \Omega^*(M)$ verifies $\|\omega\|_S \leq 0$ for any singular stratum $S \in \mathbf{S}_{\mathcal{F}}$ (cf. 2.1).

²We refer the reader to [10],[11] for details.

³Through the restriction $\omega \mapsto \omega_{R_{\mathcal{F}}}$.

⁴The map $P_{\varphi}: \mathbb{R}^{m-n-1} \times \mathbb{S}^n \times [0, 1] \rightarrow U$ is defined by $P_{\varphi}(x, y, t) = \varphi(x, [y, t])$.

2.3 Basic cohomology. The basic cohomology of the foliation \mathcal{F} is an important tool to study its transversal structure and plays the rôle of the cohomology of the orbit space M/\mathcal{F} , which can be a wild topological espace. A differential form $\omega \in \Omega^*(M)$ is *basic* if $i_X\omega = i_Xd\omega = 0$, for each vector field X on M tangent to the foliation \mathcal{F} . For exemple, a function f is basic iff f is constant on the leaves of \mathcal{F} . We shall write $\Omega^*(M/\mathcal{F})$ for the complex of basic forms. Its cohomology $H^*(M/\mathcal{F})$ is the *basic cohomology* of (M, \mathcal{F}) . We also use the *relative basic cohomology* $H^*((M, F)/\mathcal{F})$, that is, the cohomology computed from the complex of basic forms vanishing on the saturated set $F \subset M$. The basic cohomology does not use the stratification $S_{\mathcal{F}}$.

2.4 Basic intersection cohomology. A *perversity* is a map $\bar{p}: S_{\mathcal{F}}^{\sigma} \rightarrow \mathbb{Z} \cup \{-\infty, \infty\}$, where $S_{\mathcal{F}}^{\sigma}$ is the family of singular strata. The *constant perversity* $\bar{\iota}$ is defined by $\bar{\iota}(S) = \iota$, where $\iota \in \mathbb{Z} \cup \{-\infty, \infty\}$.

The basic intersection cohomology appears when one considers basic perverse forms whose perverse degree is controlled by a perversity. We shall put

$$\Omega_{\bar{p}}^*(M/\mathcal{F}) = \left\{ \omega \in \Pi_{\mathcal{F}}^*(M) / \omega \text{ is basic and } \max(\|\omega\|_s, \|d\omega\|_s) \leq \bar{p}(S) \quad \forall S \in S_{\mathcal{F}}^{\sigma} \right\}$$

the complex of basic perverse forms whose perverse degree (and that of the their derivative) is bounded by the perversity \bar{p} . The cohomology $H_{\bar{p}}^*(M/\mathcal{F})$ of this complex is the *basic intersection cohomology*⁵ of (M, \mathcal{F}) relatively to the perversity \bar{p} .

Consider a twisted product $K \times_H N$. Perversities on $K \times_H N$ and $K \times N$ are determinate by perversities on N by the formula (cf. 1.3(b)):

$$(1) \quad \bar{p}(K \times S) = \bar{p}(\Pi(K \times S)) = \bar{p}(S).$$

2.5 Mayer-Vietoris. This is the technique we use in order to decompose the manifold in nicer pieces. An open covering $\{U, V\}$ of M by saturated open subsets is a *basic covering*. It possesses a subordinated partition of the unity made up of basic functions defined on M (see [9]). For a such covering we have the Mayer-Vietoris short sequence

$$0 \rightarrow \Omega_{\bar{p}}^*(M/\mathcal{F}) \rightarrow \Omega_{\bar{p}}^*(U/\mathcal{F}) \oplus \Omega_{\bar{p}}^*(V/\mathcal{F}) \rightarrow \Omega_{\bar{p}}^*((U \cap V)/\mathcal{F}) \rightarrow 0,$$

where the map are defined by $\omega \mapsto (\omega, \omega)$ and $(\alpha, \beta) \mapsto \alpha - \beta$. The third map is onto since the elements of the partition of the unity are *controlled functions*, id est, elements of $\Omega_{\bar{p}}^0(-)$ (cf. 2.2). Thus, the sequence is exact. This result is not longer true for more general coverings.

We shall use in this work the two following local calculations (see [11, Proposition 3.5.1 and Proposition 3.5.2] for the proofs).

Proposition 2.6 *Let $(\mathbb{R}^k, \mathcal{H})$ be a simple foliation. Consider \bar{p} a perversity on M and define the perversity \bar{p} on $\mathbb{R}^k \times M$ by $\bar{p}(\mathbb{R}^k \times S) = \bar{p}(S)$. The canonical projection $\text{pr}: \mathbb{R}^k \times M \rightarrow M$ induces the isomorphism*

$$H_{\bar{p}}^*(M/\mathcal{F}) \cong H_{\bar{p}}^*(\mathbb{R}^k \times M/\mathcal{H} \times \mathcal{F}).$$

Proposition 2.7 *Let \mathcal{G} be a conical foliation without 0-dimensional leaves on the sphere \mathbb{S}^n . A perversity \bar{p} on $c\mathbb{S}^n$ gives the perversity \bar{p} on \mathbb{S}^n defined by $\bar{p}(S) = \bar{p}(S \times]0, 1[)$. The canonical projection $\text{pr}: \mathbb{S}^n \times]0, 1[\rightarrow \mathbb{S}^n$ induces the isomorphism*

$$H_{\bar{p}}^i(c\mathbb{S}^n/c\mathcal{G}) = \begin{cases} H_{\bar{p}}^i(\mathbb{S}^n/\mathcal{G}) & \text{if } i \leq \bar{p}(\{\vartheta\}) \\ 0 & \text{if } i > \bar{p}(\{\vartheta\}). \end{cases}$$

⁵BIC for short.

In the next section we shall need the following technical Lemma.

Lemma 2.8 *Let $\Phi: K \times M \rightarrow M$ be a smooth action, where K is a compact Lie group, and let V be a fundamental vector field of this action. Consider a normal subgroup G of K and write \mathcal{F} the associated conical foliation on M . Then, the interior operator $i_V: \Omega_{\bar{p}}^*(M/\mathcal{F}) \rightarrow \Omega_{\bar{p}}^{*-1}(M/\mathcal{F})$ is well defined, for any perversity \bar{p} .*

Proof. Since the question is a local one, then it suffices to consider where M is a twisted product $K \times_H N$ ⁶. Notice that the blow up $\Pi: K \times N \rightarrow K \times_H N$ is a K -equivariant map relatively to the action $\ell \cdot (k, z) = (\ell \cdot k, z)$. This gives $\Pi_*(X^u, 0) = V$ for some $u \in \mathfrak{k}$. From Lemma 3.1 we know that it suffices to prove that the operator

$$i_{(X^u, 0)}: \Omega_{\bar{p}}^*(K \times N/\mathcal{K} \times \mathcal{N}) \rightarrow \Omega_{\bar{p}}^{*-1}(K \times N/\mathcal{K} \times \mathcal{N})$$

is well defined. Since $G \triangleleft K$ then the vector field X^u preserves the foliation \mathcal{K} . So, it suffices to prove that the operator

$$i_{(X^u, 0)}: \Omega_{\bar{p}}^*(K \times N) \rightarrow \Omega_{\bar{p}}^{*-1}(K \times N)$$

is well defined. This comes from the fact that X^u acts on the K -factor while the perversion conditions are measured on the N -factor (cf. (1)). ♣

3 The BIC of a twisted product

We compute now the BIC of a twisted product $K \times_H N$ (cf. 1.3) for a perversity \bar{p} (cf. (1)).

Lemma 3.1 *The natural projection $\Pi: K \times N \rightarrow K \times_H N$ induces the differential monomorphism*

$$(2) \quad \Pi^*: \Omega_{\bar{p}}^*(K \times_H N/\mathcal{W}) \rightarrow \Omega_{\bar{p}}^*(K \times N/\mathcal{K} \times \mathcal{N}).$$

Moreover, given a differential form ω on $K \times_H R_{\mathcal{W}}$, we have:

$$(3) \quad \Pi^* \omega \in \Omega_{\bar{p}}^*(K \times N/\mathcal{K} \times \mathcal{N}) \iff \omega \in \Omega_{\bar{p}}^*(K \times_H N/\mathcal{W}).$$

Proof. Notice that the injectivity of Π^* comes from the fact that Π is a surjection. For the rest, we proceed in several steps.

(a) A foliated atlas for $\pi: K \rightarrow K/H$.

Since $\pi: K \rightarrow K/H$ is a H -principal bundle then it possesses an atlas $\mathcal{A} = \{\varphi: \pi^{-1}(U) \rightarrow U \times H\}$ made up with H -equivariant charts: $\varphi(k \cdot h^{-1}) = (\pi(k), h \cdot h_0)$ if $\varphi(k) = (\pi(k), h_0)$. We study the foliation $\varphi_* \mathcal{K}$. This equivariance property gives $\varphi_* X_u = (0, Z^u)$ for each $u \in \mathfrak{g} \cap \mathfrak{h}$. Thus, the trace of the foliation $\varphi_* \mathcal{K}$ on the fibers of the canonical projection $\text{pr}: U \times H \rightarrow U$ is C . On the other hand, since the map π is a G -equivariant map then $\pi_* \mathcal{K} = \mathcal{D}$, which gives $\text{pr}_* \varphi_* \mathcal{K} = \mathcal{D}$. We conclude that $\varphi_* \mathcal{K} \subset \mathcal{D} \times C$. By dimension reasons we get $\varphi_* \mathcal{K} = \mathcal{D} \times C$. The atlas \mathcal{A} is an H -equivariant foliated atlas of π .

(b) A foliated atlas for $\Pi: K \times N \rightarrow K \times_H N$.

⁶In fact, N is an euclidean space \mathbb{R}^a et Θ is an orthogonal action.

We claim that $\mathcal{A}_\# = \{\bar{\varphi}: \pi^{-1}(U) \times_H N \rightarrow U \times N / (U, \varphi) \in \mathcal{A}\}$ is a foliated atlas of $K \times_H N$ where the map $\bar{\varphi}$ is defined by $\bar{\varphi}(< k, z >) = (\pi(k), (\Theta((\varphi^{-1}(\pi(k), e))^{-1} \cdot k, z)))$. This map is a diffeomorphism whose inverse is $\bar{\varphi}^{-1}(u, z) = < \varphi^{-1}(u, e), z >$. It verifies

$$\bar{\varphi}_* \mathcal{W} \stackrel{1.3(a)}{=} \bar{\varphi}_* \Pi_*(\mathcal{K} \times I) = \bar{\varphi}_* \Pi_*(\varphi^{-1} \times \mathbb{I}_N)_*(\mathcal{D} \times C \times I).$$

A straightforward calculation shows $\bar{\varphi}_* \Pi_*(\varphi^{-1} \times \mathbb{I}_N) = (\mathbb{I}_U \times \Theta)$. Since C is defined by the action Γ then $\Theta_*(C \times I) = \mathcal{N}$. Finally we obtain $\bar{\varphi}_* \mathcal{W} = \mathcal{D} \times \mathcal{N}$.

(c) Last Step.

Given $(U, \varphi) \in \mathcal{A}_\#$, we have the commutative diagram

$$\begin{array}{ccc} U \times H \times N & \xrightarrow{\varphi^{-1} \times \mathbb{I}_N} & K \times N \\ \downarrow Q & & \downarrow \Pi \\ U \times N & \xrightarrow{\bar{\varphi}^{-1}} & K \times_H N \end{array}$$

where $Q(u, h, z) = (u, h^{-1} \cdot z)$, $\Pi^{-1}(\text{Im } \bar{\varphi}^{-1}) = \text{Im } (\varphi^{-1} \times \mathbb{I}_N)$ and the rows are foliated imbeddings. Now, since (2) and (3) are local questions then it suffices to prove that

- $Q^*: \Omega_p^*(U \times N / \mathcal{D} \times \mathcal{N}) \rightarrow \Omega_p^*(U \times H \times N / \mathcal{D} \times C \times \mathcal{N})$ is well-defined, and
- $Q^* \omega \in \Omega_p^*(U \times H \times N / \mathcal{D} \times C \times \mathcal{N}) \iff \omega \in \Omega_p^*(U \times N / \mathcal{D} \times \mathcal{N})$, for any $\omega \in \Omega^*(U \times R_N)$.

This comes from the fact that the map

$$\nabla: (U \times H \times N, \mathcal{D} \times C \times \mathcal{N}) \rightarrow (U \times H \times N, \mathcal{D} \times C \times \mathcal{N}),$$

defined by $\nabla(u, h, z) = (u, h, h^{-1} \cdot z)$, is a foliated diffeomorphism⁷ and $Q = \text{pr}_0 \circ \nabla$, with $\text{pr}_0: U \times H \times N \rightarrow U \times N$ canonical projection (cf. Proposition 2.6). ♣

3.2 The Lie algebra \mathfrak{k} . We suppose in this paragraph that G is also dense on K . Choose ν a bi-invariant riemannian metric on K , which exists by compactness. Consider

$$B = \{u_1, \dots, u_a, u_{a+1}, \dots, u_b, u_{b+1}, \dots, u_c, u_{c+1}, \dots, u_f\}$$

an orthonormal basis of the Lie algebra \mathfrak{k} of K with $\{u_1, \dots, u_b\}$ basis of the Lie algebra \mathfrak{g} of G and $\{u_{a+1}, \dots, u_c\}$ basis of the Lie algebra \mathfrak{h} of H . For each indice $1 \leq i \leq f$ we shall write $X_i \equiv X_{u_i}$ and $X^i \equiv X^{u_i}$ (cf. 1.2).

Let $\gamma_i \in \Omega^1(K)$ be the dual form of X_i , that is, $\gamma_i = i_{X_i} \nu$. Notice that $\delta_{ij} = \gamma_j(X_i)$. These forms are invariant by the left action of K . Since the flow of X^j is the multiplication on the left by $\exp(tu_j)$ then $L_{X^j} \gamma_i = 0$ for each $1 \leq j \leq f$.

For the differential, we have the formula $d\gamma_l = \sum_{1 \leq i < j \leq f} C_{ij}^l \gamma_i \wedge \gamma_j$, where $[X_i, X_j] = \sum_{l=1}^f C_{ij}^l X_l$, and $1 \leq i, j, l \leq f$. We have several restrictions on these coefficients. Since $G \triangleleft K$ then \mathfrak{g} is an ideal of \mathfrak{k} and therefore we have

$$C_{ij}^l = 0 \text{ for } i \leq b < l.$$

⁷Since $G \cap H \triangleleft H$.

Since K/G is an abelian group (cf. Proposition 1.1.2) then the induced bracket on $\mathfrak{k}/\mathfrak{g}$ is zero and therefore we have

$$C_{ij}^l = 0 \text{ for } b < i, j, l \leq f.$$

These equations imply that

$$(4) \quad d\gamma_l = 0 \text{ for each } b < l.$$

The \mathcal{E} -basic differential forms in $\bigwedge^* (\gamma_1, \dots, \gamma_f)$ are exactly $\bigwedge^* (\gamma_{c+1}, \dots, \gamma_f)$ since they are cycles and the family $\{X_1, \dots, X_c\}$ generates the foliation \mathcal{E} . This gives

$$(5) \quad H^*(K/\mathcal{E}) = \bigwedge^* (\gamma_{c+1}, \dots, \gamma_f).$$

3.3 Two actions of H/H_0 . The Lie group H preserves the foliation \mathcal{N} since the Lie group $G \cap H$ is a normal subgroup of H . Put H_0 the connected component of H containing the unity element. Since it is a connected compact Lie group then a standard argument shows that

$$(6) \quad \left(H_{\overline{p}}^*(N/\mathcal{N})\right)^{H_0} = H^*\left(\left(\Omega_{\overline{p}}^*(N/\mathcal{N})\right)^{H_0}\right) = H_{\overline{p}}^*(N/\mathcal{N})$$

(cf. [5, Theorem I, Ch. IV, vol. II]). We conclude that the finite group H/H_0 acts naturally on $H_{\overline{p}}^*(N/\mathcal{N})$.

Since H_0 is a connected Lie subgroup of GH then $\left(H^*(K/\mathcal{E})\right)^{H_0} = H^*(K/\mathcal{E})$. We conclude that the finite group H/H_0 acts naturally on $H^*(K/\mathcal{E})$.

Proposition 3.4 *Let (K, G, H) be a trio with G connected and dense in K . Then*

$$H_{\overline{p}}^*(K \times_H N/\mathcal{W}) = \left(H^*(K/\mathcal{E}) \otimes H_{\overline{p}}^*(N/\mathcal{N})\right)^{H/H_0}.$$

Proof. Using the blow up $\Pi: K \times N \rightarrow K \times_H N$, the computation of $H_{\overline{p}}^*(K \times_H N/\mathcal{W})$ can be done by using the complex $\text{Im} \left\{ \Pi^*: \Omega_{\overline{p}}^*(K \times_H N/\mathcal{F}) \rightarrow \Omega_{\overline{p}}^*(K \times N/\mathcal{K} \times \mathcal{N}) \right\}$ (cf. Lemma 3.1). We study this complex in several steps. We fix $B = \{u_1, \dots, u_f\}$ an orthonormal basis of \mathfrak{k} as in 3.2.

i) Description of $\Omega^*(K \times R_{\mathcal{N}})$.

A differential form $\omega \in \Omega^*(K \times R_{\mathcal{N}})$ is of the form

$$(7) \quad \eta + \sum_{1 \leq i_1 < \dots < i_\ell \leq f} \gamma_{i_1} \wedge \dots \wedge \gamma_{i_\ell} \wedge \eta_{i_1, \dots, i_\ell},$$

where the forms $\eta, \eta_{i_1, \dots, i_\ell} \in \Omega^*(K \times R_{\mathcal{N}})$ verify $i_{X_j} \eta = i_{X_j} \eta_{i_1, \dots, i_\ell} = 0$ for each $1 \leq j \leq f$ and each $1 \leq i_1 < \dots < i_\ell \leq f$.

ii) Description of $\Pi_{\mathcal{K} \times \mathcal{N}}^*(K \times N)$.

Since the foliation \mathcal{K} is regular then we always can choose a conical chart of the form $(U_1 \times U_2, \varphi_1 \times \varphi_2)$ where (U_1, φ_1) is a foliated chart of (K, \mathcal{K}) and (U_2, φ_2) is a conical chart of (N, \mathcal{N}) . The local blow up of the chart $(U_1 \times U_2, \varphi_1 \times \varphi_2)$ is constructed from the second factor without modifying the first one. So, the differential forms γ_i are always perverse forms and a differential form $\omega \in \Pi_{\mathcal{K} \times \mathcal{N}}^*(K \times N)$ is of the form (7) where $\eta, \eta_{i_1, \dots, i_\ell} \in \Pi_{\mathcal{K} \times \mathcal{N}}^*(K \times N)$ verify $i_{X_j} \eta = i_{X_j} \eta_{i_1, \dots, i_\ell} = 0$ for each $1 \leq j \leq f$ and each $1 \leq i_1 < \dots < i_\ell \leq f$.

⟨iii⟩ Description of $\Omega^*(K \times R_N/\mathcal{K} \times \mathcal{N})$.

Take $\omega \in \Omega^*(K \times R_N/\mathcal{K} \times \mathcal{N})$. Since \mathcal{K} is generated by the family $\{X_j / 1 \leq j \leq b\}$ then $L_{X_j}\omega = 0$ for any $1 \leq j \leq b$, or equivalently, $R_g^*\omega = \omega$ for each $g \in G$ since G is connected. By density, $R_k^*\omega = \omega$ for each $k \in K$ and therefore $L_{X_j}\omega = 0$ for any $1 \leq j \leq f$ since K is connected. We conclude that $L_{X_j}\eta = L_{X_j}\eta_{i_1, \dots, i_\ell} = 0$ for any $1 \leq j \leq f$ and each $1 \leq i_1 < \dots < i_\ell \leq f$. This gives $\omega \in \bigwedge^*(\gamma_1, \dots, \gamma_f) \otimes \Omega^*(R_N)$.

The \mathcal{N} -basic differential forms of $\Omega^*(R_N)$ are exactly $\Omega^*(R_N/\mathcal{N})$. The \mathcal{K} -basic differential forms of $\bigwedge^*(\gamma_1, \dots, \gamma_f)$ are exactly $\bigwedge^*(\gamma_{b+1}, \dots, \gamma_f)$ (cf. (4)). From these two facts, we get

$$\Omega^*(K \times R_N/\mathcal{K} \times \mathcal{N}) = \bigwedge^*(\gamma_{b+1}, \dots, \gamma_f) \otimes \Omega^*(R_N/\mathcal{N})$$

as differential graduate commutative algebras.

⟨iv⟩ Description of $\Omega_p^*(K \times N/\mathcal{K} \times \mathcal{N})$.

From ⟨ii⟩ and ⟨iii⟩ it suffices to control the perverse degree of the forms

$$\eta + \sum_{b+1 \leq i_1 < \dots < i_\ell \leq f} \gamma_{i_1} \wedge \dots \wedge \gamma_{i_\ell} \wedge \eta_{i_1, \dots, i_\ell} \in \bigwedge^*(\gamma_{b+1}, \dots, \gamma_f) \otimes \Pi_N^*(N).$$

Consider S a stratum of \mathbf{S}_N . From $\|\gamma_i\|_{K \times S} = 0$ and $\|\eta\|_{K \times S} = \|\eta\|_S$, we get $\|\gamma_{i_1} \wedge \dots \wedge \gamma_{i_\ell} \wedge \eta_{i_1, \dots, i_\ell}\|_{K \times S} = \|\eta_{i_1, \dots, i_\ell}\|_S$. We conclude that

$$\Omega_p^*(K \times N/\mathcal{K} \times \mathcal{N}) \cong \bigwedge^*(\gamma_{b+1}, \dots, \gamma_f) \otimes \Omega_p^*(N/\mathcal{N})$$

(cf. 1.3(b)).

⟨v⟩ Description of $\text{Im } \{\Pi^*: \Omega_p^*(K \times_H N/\mathcal{F}) \longrightarrow \Omega_p^*(K \times N/\mathcal{K} \times \mathcal{N})\}$.

We denote by $\{W_{a+1}, \dots, W_c\}$ the fundamental vector fields of the action $\Theta: H \times N \rightarrow N$ associated to the basis $\{u_{a+1}, \dots, u_c\}$. Consider now the action $\Upsilon: H \times (K \times N) \rightarrow (K \times N)$ defined by $\Upsilon(h, (k, z)) = (k \cdot h^{-1}, \Theta(h, z))$. Its fundamental vector fields associated to the basis $\{u_{a+1}, \dots, u_c\}$ are $\{(X_{a+1}, W_{a+1}), \dots, (X_c, W_c)\}$. Given $h \in H$, we take $\Upsilon_h: K \times N \rightarrow K \times N$ the map defined by $\Upsilon_h(k, z) = \Upsilon(h, (k, z))$. Then, we have

$$\text{Im } \Pi^* = \left\{ \omega \in \bigwedge^*(\gamma_{b+1}, \dots, \gamma_f) \otimes \Omega_p^*(N/\mathcal{N}) \left| \begin{array}{l} \text{(i) } i_{X_i}\omega = -i_{W_i}\omega \text{ if } a < i \leq c \\ \text{(ii) } L_{X_i}\omega = -L_{W_i}\omega \text{ if } a < i \leq c, \\ \text{(iii) } (\Upsilon_h)^*\omega = \omega \text{ for } h \in H. \end{array} \right. \right\}.$$

Let H_0 be the unity connected component of H . Recall that the subgroup H_0 is normal in H and that the quotient H/H_0 is a finite group. Conditions (ii) gives that ω is H_0 -invariant. So, condition (iii) can be replaced by: (iv) $(\Upsilon_h)^*\omega = \omega$ for $h \in H/H_0$. Therefore

$$\text{Im } \Pi^* = \left\{ \omega \in \bigwedge^*(\gamma_{b+1}, \dots, \gamma_f) \otimes \Omega_p^*(N/\mathcal{N}) \left| \begin{array}{l} \text{(i) } i_{X_i}\omega = -i_{W_i}\omega \text{ if } a < i \leq c \\ \text{(ii) } L_{X_i}\omega = -L_{W_i}\omega \text{ if } a < i \leq c. \end{array} \right. \right\}^{H/H_0}.$$

Since the group H/H_0 is a finite one, we get that the cohomology $H^*(\text{Im } \Pi^*)$ is isomorphic to $(H^*(A))^{\cdot H/H_0}$, where A^* is the differential complex

$$\left\{ \omega \in \bigwedge^*(\gamma_{b+1}, \dots, \gamma_f) \otimes \Omega_p^*(N/\mathcal{N}) \left| \begin{array}{l} \text{(i) } i_{X_i}\omega = -i_{W_i}\omega \text{ if } a < i \leq c \\ \text{(ii) } L_{X_i}\omega = -L_{W_i}\omega \text{ if } a < i \leq c \end{array} \right. \right\}.$$

So, it remains to compute $H^*(A)$. This computation can be simplified by using these three facts:

- $i_{W_i}\omega = L_{W_i}\omega = 0$ for each $a < i \leq b$, since the foliation \mathcal{N} is defined by the action of $G \cap H$.
- $i_{X_i}\gamma_j = \delta_{ij}$ for all i, j (cf. 3.2).
- $d\gamma_j = 0$ for $b < j$ (cf. (4)).

We get that A^* is the differential complex

$$\left\{ \omega \in \bigwedge^* (\gamma_{b+1}, \dots, \gamma_f) \otimes \Omega_{\overline{p}}^*(N/\mathcal{N}) \left| \begin{array}{l} \text{(i) } i_{X_i}\omega = -i_{W_i}\omega \text{ if } b < i \leq c \\ \text{(ii) } 0 = L_{W_i}\omega \text{ if } b < i \leq c \end{array} \right. \right\} =$$

$$\bigwedge^* (\gamma_{c+1}, \dots, \gamma_f) \otimes \underbrace{\left\{ \omega \in \bigwedge^* (\gamma_{b+1}, \dots, \gamma_c) \otimes \Omega_{\overline{p}}^*(N/\mathcal{N}) \left| \begin{array}{l} \text{(i) } i_{X_i}\omega = -i_{W_i}\omega \text{ if } b < i \leq c \\ \text{(ii) } 0 = L_{W_i}\omega \text{ if } b < i \leq c \end{array} \right. \right\}}_{B^*}.$$

A straightforward computation gives that the canonical writing of a form $\omega \in \bigwedge^* (\gamma_{b+1}, \dots, \gamma_c) \otimes \Omega_{\overline{p}}^*(N/\mathcal{N})$ verifying (i) is

$$(8) \quad \omega = \omega_0 + \sum_{b < i_1 < \dots < i_\ell \leq c} (-1)^\ell \gamma_{i_1} \wedge \dots \wedge \gamma_{i_\ell} \wedge (i_{W_{i_\ell}} \dots i_{W_{i_1}} \omega_0)$$

for some $\omega_0 \in \Omega_{\overline{p}}^*(N/\mathcal{N})$ (cf. Lemma 2.8).

Consider now $b < i, j \leq c$. Since K/G is an abelian group (cf. Proposition 1.1.2) and H is a Lie group then $[W_i, W_j] = \sum_{l=a+1}^b C_{ij}^l W_l$. Then, $i_{[W_i, W_j]}\omega_0 = 0$ since the foliation \mathcal{N} is defined by the action of $G \cap H$. So, the canonical writing of a form $\omega \in B^*$ is (8) for some $\omega_0 \in \{\eta \in \Omega_{\overline{p}}^*(N/\mathcal{N}) / L_{W_i}\eta = 0 \text{ if } b < i \leq c\} = (\Omega_{\overline{p}}^*(N/\mathcal{N}))^{H_0}$.

Then, the operator $\Delta: B^* \longrightarrow (\Omega_{\overline{p}}^*(N/\mathcal{N}))^{H_0}$, defined by $\Delta(\omega) = \omega_0$, is a differential isomorphism. We conclude that the differential complex A^* is isomorphic to $\bigwedge^* (\gamma_{c+1}, \dots, \gamma_f) \otimes (\Omega_{\overline{p}}^*(N/\mathcal{N}))^{H_0}$ and therefore $H^*(A) \cong H^*(K/\mathcal{E}) \otimes H_{\overline{p}}^*(N/\mathcal{N})$ (cf. (5) and (6)). Since the operator Δ is (H/H_0) -equivariant (cf. 3.3) then we get

$$H_{\overline{p}}^*(K \times_H N/\mathcal{W}) = H^*(\text{Im } \Pi^*) = (H^*(A))^{H/H_0} = (H^*(K/\mathcal{E}) \otimes H_{\overline{p}}^*(N/\mathcal{N}))^{H/H_0}.$$

This ends the proof. ♣

3.5 Remarks.

(a) When the Lie group G is commutative then K is also commutative. Differential forms γ_\bullet are K -invariants on the left and on the right, so $(H^*(K/\mathcal{E}))^H = H^*(K/\mathcal{E})$ and therefore

$$H_{\overline{p}}^*(K \times_H N/\mathcal{W}) = H^*(K/\mathcal{E}) \otimes (H_{\overline{p}}^*(N/\mathcal{N}))^{H/H_0} = H^*(K/\mathcal{E}) \otimes (H_{\overline{p}}^*(N/\mathcal{N}))^H$$

as it has been proved in [11, Proposition 3.8.4].

(b) Since the foliation \mathcal{E} is a riemannian foliation defined on a compact manifold then we know that the cohomology $H^*(K/\mathcal{E})$ is finite (cf. [4]). So, the finiteness of $H_{\overline{p}}^*(K \times_H N/\mathcal{W})$ depends on the finiteness of $H_{\overline{p}}^*(N/\mathcal{N})$.

4 Finiteness of the BIC

We prove in this section that the BIC of a Killing foliation on a compact manifold is finite dimensional. First of all, we present two geometrical tools we shall use in the proof: the isotropy type stratification and the Molino's blow up.

We fix an isometric action $\Phi: G \times M \rightarrow M$ on the compact manifold M . We denote by \mathcal{F} the induced Killing foliation. For the study of \mathcal{F} we can suppose that G is connected (see Lemma 1.1.1). We fix K a tamer group. Notice that the group G is normal in K and the quotient K/G is commutative (cf. Proposition 1.1.2).

4.1 Isotropy type stratification. The *isotropy type stratification* $\mathbf{S}_{K,M}$ of M is defined by the equivalence relation⁸:

$$x \sim y \Leftrightarrow K_x \text{ is conjugated to } K_y.$$

When $\text{depth } \mathbf{S}_{K,M} > 0$, any closed stratum $S \in \mathbf{S}_{K,M}$ is a K -invariant submanifold of M and then it possesses a K -invariant tubular neighborhood $(T, \tau, S, \mathbb{R}^m)$ whose structural group is $O(m)$. Recall that there are the following smooth maps associated with this neighborhood:

- + The *radius map* $\rho: T \rightarrow [0, 1[$ defined fiberwise from the assignation $[x, t] \mapsto t$. Each $t \neq 0$ is a regular value of the ρ . The pre-image $\rho^{-1}(0)$ is S . This map is K -invariant, that is, $\rho(k \cdot z) = \rho(z)$.
- + The *contraction* $H: T \times [0, 1] \rightarrow T$ defined fiberwisely from $([x, t], r) \mapsto [x, rt]$. The restriction $H_t: T \rightarrow T$ is an embedding for each $t \neq 0$ and $H_0 \equiv \tau$. We shall write $H(z, t) = t \cdot z$. This map is K -invariant, that is, $t \cdot (k \cdot z) = k \cdot (t \cdot z)$.

The hyper-surface $D = \rho^{-1}(1/2)$ is the *tube* of the tubular neighborhood. It is a K -invariant submanifold of T . Notice that the map

$$\nabla: D \times [0, 1[\longrightarrow T,$$

defined by $\nabla(z, t) = (2t) \cdot z$ is a K -equivariant smooth map, where K acts trivially on the $[0, 1[$ -factor. Its restriction $\nabla: D \times]0, 1[\longrightarrow T \setminus S$ is a K -equivariant diffeomorphism.

Denote S_{\min} the union of closed (minimal) strata and choose T_{\min} a disjoint family of K -invariant tubular neighborhoods of the closed strata. The union of associated tubes is denoted by D_{\min} . Notice that the induced map $\nabla_{\min}: D_{\min} \times]0, 1[\longrightarrow T_{\min} \setminus S_{\min}$ is a K -equivariant diffeomorphism.

4.2 Molino's blow up. The Molino' blow up [7] of the foliation \mathcal{F} produces a new foliation $\widehat{\mathcal{F}}$ of the same kind but of smaller depth. We suppose $\text{depth } \mathbf{S}_{K,M} > 0$. The *blow up* of M is the compact manifold

$$\widehat{M} = \left\{ (D_{\min} \times]-1, 1[) \bigsqcup ((M \setminus S_{\min}) \times \{-1, 1\}) \right\} / \sim,$$

where $(z, t) \sim (\nabla_{\min}(z, |t|), t/|t|)$, and the map $\mathcal{L}: \widehat{M} \longrightarrow M$ defined by

$$\mathcal{L}(v) = \begin{cases} \nabla_{\min}(z, |t|) & \text{if } v = (z, t) \in D_{\min} \times]-1, 1[\\ z & \text{if } v = (z, j) \in (M \setminus S_{\min}) \times \{-1, 1\}. \end{cases}$$

Notice that \mathcal{L} is a continuous map whose restriction $\mathcal{L}: \widehat{M} \setminus \mathcal{L}^{-1}(S_{\min}) \rightarrow M \setminus S_{\min}$ is a K -equivariant smooth trivial 2-covering.

⁸For notions related with compact Lie group actions, we refer the reader to [1].

Since the map ∇_{\min} is K -equivariant then Φ induces the action $\widehat{\Phi}: K \times \widehat{M} \longrightarrow \widehat{M}$ by saying that the blow-up \mathcal{L} is K -equivariant. The open submanifolds $\mathcal{L}^{-1}(T_{\min})$ and $\mathcal{L}^{-1}(T_{\min} \setminus S_{\min})$ are clearly K -diffeomorphic to $D_{\min} \times]-1, 1[$ and $D_{\min} \times (]-1, 0[\cup]0, 1[)$ respectively.

The restriction $\widehat{\Phi}: G \times \widehat{M} \longrightarrow \widehat{M}$ is an isometric action with K as a tamer group. The induced Killing foliation is $\widehat{\mathcal{F}}$. Foliations \mathcal{F} and $\widehat{\mathcal{F}}$ are related by \mathcal{L} which is a foliated map. Moreover, if S is a not minimal stratum of $\mathbf{S}_{K,M}$ then there exists a unique stratum $S' \in \mathbf{S}_{K,\widehat{M}}$ such that $\mathcal{L}^{-1}(S) \subset S'$. The family $\{S' / S \in \mathbf{S}_{K,M}\}$ covers \widehat{M} and verifies the relationship: $S_1 < S_2 \Leftrightarrow S'_1 < S'_2$. We conclude the important property

$$(9) \quad \text{depth } \mathbf{S}_{K,\widehat{M}} < \text{depth } \mathbf{S}_{K,M}.$$

4.3 Finiteness of a tubular neighborhood. We suppose $\text{depth } \mathbf{S}_{K,M} > 0$. Consider a closed stratum $S \in \mathbf{S}_{K,M}$. Take $(T, \tau, S, \mathbb{R}^m)$ a K -invariant tubular neighborhood. We fix a base point $x \in S$. The isotropy subgroup K_x acts orthogonally on the fiber $\mathbb{R}^m = \tau^{-1}(x)$. So, the induced action $\Lambda_x: G_x \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ is an isometric action, it gives the Killing foliation \mathcal{N} on \mathbb{R}^m .

Proposition 4.3.1 *If the BIC of $(\mathbb{R}^m, \mathcal{N})$ is finite dimensional then the BIC of (T, \mathcal{F}) is also finite dimensional.*

Proof. We proceed in two steps.

(a) $K_y = K_x$ for each $y \in S$.

The canonical projection $\pi: S \rightarrow S/K$ is an homogeneous bundle with fiber K/K_x . For any open subset $V \subset S/K$ the pull back $\tau^{-1}\pi^{-1}(V)$ is a K -invariant subset of T , then we can apply the Mayer-Vietoris technics to this kind of subsets (cf. 2.5).

Since the manifold S/K is a compact one then we can find a finite good covering $\{U_i / i \in I\}$ of it (cf. [2]). An inductive argument on the cardinality of I reduces the proof of the Lemma to the case where $T = \tau^{-1}\pi^{-1}(V)$, where V is a contractible open subset of S/K .

Here, the manifold T is K -equivalently diffeomorphic to $V \times (K \times_{K_x} \mathbb{R}^m)$, where K does not act on the first factor. So, the natural retraction of V to a point gives a K -equivariant retraction of T to the twisted product $K \times_{K_x} \mathbb{R}^m$. Now the result comes directly from 3.5(b) since (K, G, K_x) is a trio.

(b) General case.

The stratum S is K -equivariantly diffeomorphic to the twisted product $K \times_{N(K_x)} F$ where $N(K_x)$ is the normalizer of K_x on K and $F = S^{K_x}$. So, the tubular neighborhood T is K -equivariantly diffeomorphic to the twisted product $K \times_{N(K_x)} N$ where N is the manifold $\tau^{-1}(F)$. The previous case gives that the BIC of (N, \mathcal{F}_N) is finite dimensional. Now the result comes directly from 3.5(b) since $(K, G, N(K_x))$ is a trio. ♣

The main result of this work is the following

Theorem 4.4 *The BIC of the foliation determined by an isometric action on a compact manifold is finite dimensional.*

Proof. Let \mathcal{F} be a Killing foliation defined on a compact manifold M induced by an isometric action $\Phi: G \times M \rightarrow M$ where G is a Lie group. Without loss of generality we can suppose that the Lie group G is a connected one (cf. Lemma 1.1.1). We fix a tamer group K . We know that G is normal in K and the quotient group K/G is commutative (cf. Proposition 1.1.2).

Let us consider the following statement

$\mathfrak{A}(U, \mathcal{F}) = \text{“The BIC } H_{\bar{p}}^*(U/\mathcal{F}) \text{ is finite dimensional for each perversity } \bar{p}\text{”}$

where $U \subset M$ is a K -invariant submanifold. We prove $\mathfrak{A}(M, \mathcal{F})$ by induction on $\dim M$. The result is clear when $\dim M = 0$. We suppose $\mathfrak{A}(W, \mathcal{F})$ for any K -invariant compact submanifold W of M with $\dim W < \dim M$ and we prove $\mathfrak{A}(M, \mathcal{F})$. We proceed in several steps.

First step: 0-depth. Let us suppose $\text{depth } \mathbf{S}_{K,M} = 0$. Since $G \triangleleft K$ and K_x is conjugated to K_y , then G_x is conjugated to G_y , $\forall x, y \in M$. We get that the foliation \mathcal{F} is a (regular) riemannian foliation (cf. [7]). Its BIC is just the basic cohomology (cf. 2.3). Then $\mathfrak{A}(M, \mathcal{F})$ comes from [4].

Second step: Inside M . Let us suppose $\text{depth } \mathbf{S}_{K,M} > 0$. The family $\{M \setminus S_{\min}, T_{\min}\}$ is a basic covering of M and we get the exact sequence (cf. 2.5)

$$0 \rightarrow \Omega_{\bar{p}}^*(M/\mathcal{F}) \rightarrow \Omega_{\bar{p}}^*((M \setminus S_{\min})/\mathcal{F}) \oplus \Omega_{\bar{p}}^*(T_{\min}/\mathcal{F}) \rightarrow \Omega_{\bar{p}}^*((T_{\min} \setminus S_{\min})/\mathcal{F}) \rightarrow 0.$$

The Five Lemma gives

$$\mathfrak{A}(T_{\min} \setminus S_{\min}, \mathcal{F}), \mathfrak{A}(T_{\min}, \mathcal{F}) \text{ and } \mathfrak{A}(M \setminus S_{\min}, \mathcal{F}) \implies \mathfrak{A}(M, \mathcal{F}).$$

Since $T_{\min} \setminus S_{\min}$ is K -diffeomorphic to $D_{\min} \times]0, 1[$ (cf. (cf. 4.1)) then $\mathfrak{A}(D_{\min}, \mathcal{F}) \implies \mathfrak{A}(T_{\min} \setminus S_{\min}, \mathcal{F})$. The inequality $\dim D_{\min} < \dim M$ gives

$$\mathfrak{A}(T_{\min}, \mathcal{F}) \text{ and } \mathfrak{A}(M \setminus S_{\min}, \mathcal{F}) \implies \mathfrak{A}(M, \mathcal{F}).$$

In order to prove $\mathfrak{A}(T_{\min}, \mathcal{F})$ it suffices to prove $\mathfrak{A}(T, \mathcal{F})$ where $(T, \tau, S, \mathbb{R}^m)$ a K -invariant tubular neighborhood of closed stratum S of $\mathbf{S}_{K,M}$. Following Proposition 4.3.1 we have

$$\mathfrak{A}(\mathbb{R}^m, \mathcal{N}) \implies \mathfrak{A}(T, \mathcal{F}) \implies \mathfrak{A}(T_{\min}, \mathcal{F}).$$

Consider the orthogonal decomposition $\mathbb{R}^m = \mathbb{R}^{m_1} \times \mathbb{R}^{m_2}$, where $\mathbb{R}^{m_1} = (\mathbb{R}^m)^{G_x}$. The only fixed point of the restriction $\Lambda_x: G_x \times \mathbb{R}^{m_2} \rightarrow \mathbb{R}^{m_2}$ is the origin. So, there exists a Killing foliation⁹ \mathcal{G} on the sphere \mathbb{S}^{m_2-1} with $(\mathbb{R}^{m_1} \times \mathbb{R}^{m_2}, \mathcal{F}) = (\mathbb{R}^{m_1} \times c\mathbb{S}^{m_2-1}, \mathcal{I} \times c\mathcal{G})$. Propositions 2.6 and 2.7 give:

$$\mathfrak{A}(\mathbb{S}^{m_2-1}, \mathcal{G}) \implies \mathfrak{A}(\mathbb{R}^{m_1} \times c\mathbb{S}^{m_2-1}, \mathcal{I} \times c\mathcal{G}) \implies \mathfrak{A}(\mathbb{R}^m, \mathcal{N}).$$

Finally, since $\dim \mathbb{S}^{m_2-1} < m \leq \dim T \leq \dim M$ we have

$$(10) \quad \mathfrak{A}(M \setminus S_{\min}, \mathcal{F}) \implies \mathfrak{A}(M, \mathcal{F}).$$

Third step: Blow-up. Let us suppose $\text{depth } \mathbf{S}_{K,M} > 0$. The family $\{\mathcal{L}^{-1}(M \setminus S_{\min}), \mathcal{L}^{-1}(T_{\min})\}$ is a basic covering of \widehat{M} and we get the exact sequence (cf. 2.5)

$$0 \rightarrow \Omega_{\bar{p}}^*(\widehat{M}/\widehat{\mathcal{F}}) \rightarrow \Omega_{\bar{p}}^*(\mathcal{L}^{-1}(M \setminus S_{\min})/\widehat{\mathcal{F}}) \oplus \Omega_{\bar{p}}^*(\mathcal{L}^{-1}(T_{\min})/\widehat{\mathcal{F}}) \rightarrow \Omega_{\bar{p}}^*(\mathcal{L}^{-1}(T_{\min} \setminus S_{\min})/\widehat{\mathcal{F}}) \rightarrow 0.$$

Following 4.2 we have that

- $\mathcal{L}^{-1}(M \setminus S_{\min})$ is K -diffeomorphic to two copies of $M \setminus S_{\min}$,
- $\mathcal{L}^{-1}(T_{\min})$ is K -diffeomorphic to $D_{\min} \times]-1, 1[$,
- $\mathcal{L}^{-1}(T_{\min} \setminus S_{\min})$ is K -diffeomorphic to $D_{\min} \times (]-1, 0[\cup]0, 1[)$.

⁹It is given by the orthogonal action $\Lambda_x: G_x \times \mathbb{S}^{m_2-1} \rightarrow \mathbb{S}^{m_2-1}$.

Now, the Five Lemma gives

$$\mathfrak{A}(D_{\min}, \widehat{\mathcal{F}}) \text{ and } \mathfrak{A}(\widehat{M}, \widehat{\mathcal{F}}) \implies \mathfrak{A}(M \setminus S_{\min}, \mathcal{F}).$$

But, the inequality $\dim D_{\min} < \dim M$ gives

$$(11) \quad \mathfrak{A}(\widehat{M}, \widehat{\mathcal{F}}) \implies \mathfrak{A}(M \setminus S_{\min}, \mathcal{F}).$$

Forth step: Final blow-up. When $\text{depth } S_{K,M} = 0$ we get $\mathfrak{A}(M, \mathcal{F})$ from the First step. Let us suppose $\text{depth } S_{K,M} > 0$. From (10) and (11) we get

$$\mathfrak{A}(\widehat{M}, \widehat{\mathcal{F}}) \implies \mathfrak{A}(M, \mathcal{F}).$$

with $\text{depth } S_{K,\widehat{M}} < \text{depth } S_{K,M}$ (cf. (9)). By iterating this procedure we get

$$\mathfrak{A}(\widehat{M}, \widehat{\mathcal{F}}) = \mathfrak{A}\left(\widehat{\widehat{M}}, \widehat{\widehat{\mathcal{F}}}\right) \implies \dots \implies \mathfrak{A}(\widehat{M}, \widehat{\mathcal{F}}) \implies \mathfrak{A}(M, \mathcal{F}),$$

with $\text{depth } S_{K,\widehat{M}} = 0$. We finish the proof by applying again the First Step. ♣

References

- [1] G. E. Bredon: *Introduction to Compact Transformation Groups* - Pure Appl. Math. **46**. Academic Press. 1972.
- [2] G. E. Bredon: *Topology and Geometry* - Graduate Text in Math. **139**. Springer. 1993.
- [3] A. El Kacimi and G. Hector *Décomposition de Hodge basique pour un feuilletage riemannien*. - Ann. Inst. Fourier **36**(1988), 207-227.
- [4] A. El Kacimi, V. Sergiescu and G. Hector *La cohomologie basique d'un feuilletage riemannien est de dimension finie* - Math. Z. **188**(1985), 593-599.
- [5] W. Greub, S. Halperin and R. Vanstone: *Connections, curvature and cohomology* - Pure and Appl. Math., vol. II, Academic Press, New York and London, 1972.
- [6] S. Kobayashi: *Transformation groups in differential geometry*. Classic in Mathematics. Springer, 1995.
- [7] P. Molino: *Riemannian Foliations* - Progress in Math. **73**. Birkhäuser, 1988.
- [8] D. Poguntke: *Dense Lie group homomorphisms* - J. Algebra **169**(1994), 625-647.
- [9] J.I. Royo Prieto, M. Saralegi-Aranguren and R. Wolak: *Tautness for riemannian foliations on non-compact manifolds*. - Manus. Math. **126** (2008) 177-200.
- [10] M. Saralegi-Aranguren and R. Wolak: *The BIC of a conical fibration*. - Mat. Zametki **77**(2005), 235–257. Translation in Math. Notes **77**(2005), 213-231.
- [11] M. Saralegi-Aranguren and R. Wolak: *The BIC of a singular foliation defined by an abelian group of isometries*. - Annal. Pol. Math. **89** (2006) 203-246.